

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Keep away from humans, domestic animals, and pets.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All handlers (including applicators) must wear long-sleeved shirt, long pants, shoes, socks, and waterproof gloves. Any person who retrieves carcasses or unused bait following application of this product must wear waterproof gloves.

USER SAFETY REQUIREMENTS

Follow manufacturer's instruction for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to mammals and birds. Dogs, cats and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store only in original closed container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals. Sweep up spillage carefully and dispose of as indicated below.

"PESTICIDE DISPOSAL"

Wastes resulting from the use of this product may be disposed of at on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container

DO NOT REUSE OR REFILL THIS CONTAINER
[Plastic] Offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

KEEP AWAY FROM FEED AND FOODSTUFFS


WILCO DISTRIBUTORS, INC.
P O BOX 291 • LOMPOC CA 93438
PH# 805-735-2476

NET WT. 4 LBS.

EPA ESTB. NO.: 36029-CA-01
EPA REG NO.: 36029-20

FPO

11523



**GROUND
SQUIRREL
BAIT
BY
WILCO**

This product may only be used within 50 feet of buildings to control California and Townsend Ground Squirrels in California, Nevada, Oregon, Washington, Idaho, and Utah only. Ground Squirrel Bait by Wilco is actually impregnated by the toxicant.

Active ingredient:

Diphacinone (CAS# 82666)..... 0.005%

Other ingredients.....99.995%

Total.....100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION
FIRST AID**

Have label with you when obtaining treatment advice.

If swallowed:	<ul style="list-style-type: none"> • Call a poison control center, doctor or 1-800-858-7378 immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor immediately for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor immediately for treatment advice.

TREATMENT FOR PET POISONING

If animal eats bait, call a veterinarian or 1-800-222-1222 at once.

NOTE TO PHYSICIAN OR VETERINARIAN

Contains the anticoagulant Diphacinone. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. For humans or animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give vitamin K1 intramuscularly or orally.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions. Use only for the sites, pets, and application directions described on this label.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help prevent accidents.

1. Store product not in use in a location out of reach of children, pets, and livestock.

2. Apply this product in secured, tamper-resistant bait stations, (like Wilco Ground Squirrel Bait Stations). These stations must be resistant to destruction by dogs and by children under six years of age, and must be constructed and used in ways which prevent such children from reaching into bait compartments and obtaining bait. Stations must be secured so that they cannot be overturned by young children, pets, or winds. Stations used in areas open to livestock and/or nontarget wildlife must be secured so that the units cannot be readily overturned by any nontarget animals that have access to them.

3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: For use only around buildings (including yards and flower gardens) in secured, tamper-resistant bait stations (like Wilco Ground Squirrel Bait Stations) to control the California Ground Squirrel (*Spermophilus beecheyi*) and Townsend Ground Squirrel (*Spermophilus townsendi*) in California, Nevada, Oregon, Washington, Idaho, and Utah only. Do not apply this product by any method not specified on this label. This product may not be broadcast. Do not sell this product in individual containers holding less than 4 pounds of bait.

APPLICATION DIRECTIONS: Secure bait stations at locations where they will be readily accessible to target ground squirrels, preferably near active burrows. Secure stations 20 to 100 feet apart, depending upon density of the population. Wear waterproof gloves and scooping device (preferably long-handled) to transfer bait from the container to the station. Use 2 to 4 pounds of bait per bait station. Check bait stations frequently to insure that an uninterrupted supply of bait is always available for at least 15 days or until signs of feeding have stopped.

POST-APPLICATION DIRECTIONS:

Wearing waterproof gloves, collect and bury dead rodents. Dispose of leftover bait according to "Pesticide Disposal". With detergent and hot water, wash all implements used for applying bait. Do not use these implements for mixing, holding, or transferring foods or feed.